

Clinical paper

Receiving hospital characteristics associated with survival after out-of-hospital cardiac arrest^{☆,☆☆}

Clifton W. Callaway^{a,*}, Robert Schmicker^b, Mitch Kampmeyer^a, Judy Powell^b,
Tom D. Rea^b, Mohamud R. Daya^c, Thomas P. Aufderheide^d, Daniel P. Davis^e,
Jon C. Rittenberger^a, Ahamed H. Idris^f, Graham Nichol^b,
The Resuscitation Outcomes Consortium (ROC) Investigators

^a University of Pittsburgh, Pittsburgh, PA, United States

^b University of Washington, Seattle, WA, United States

^c Oregon Health and Science University, Portland, OR, United States

^d Medical College of Wisconsin, Milwaukee, WI, United States

^e University of California, San Diego, San Diego, CA, United States

^f University of Texas, Southwestern Medical Center, Dallas, TX, United States

ARTICLE INFO

Article history:

Received 14 October 2009

Received in revised form

30 November 2009

Accepted 3 December 2009

Keywords:

Heart arrest

Regionalization

Catheterization

Post-resuscitation care

ABSTRACT

Aim: Survival after out-of-hospital cardiac arrest (OOHCA) varies between regions, but the contribution of different factors to this variability is unknown. This study examined whether survival to hospital discharge was related to receiving hospital characteristics, including bed number, capability of performing cardiac catheterization and hospital volume of OOHCA cases.

Material and methods: Prospective observational database of non-traumatic OOHCA assessed by emergency medical services was created in 8 US and 2 Canadian sites from December 1, 2005 to July 1, 2007. Subjects received hospital care after OOHCA, defined as either (1) arriving at hospital with pulses, or (2) arriving at hospital without pulses, but discharged or died ≥ 1 day later.

Results: A total of 4087 OOHCA subjects were treated at 254 hospitals, and 32% survived to hospital discharge. A majority of subjects (68%) were treated at 116 (46%) hospitals capable of cardiac catheterization. Unadjusted survival to discharge was greater in hospitals performing cardiac catheterization (34% vs. 27%, $p = 0.001$), and in hospitals that received ≥ 40 patients/year compared to those that received < 40 (37% vs. 30%, $p = 0.01$). Survival was not associated with hospital bed number, teaching status or trauma center designation. Length of stay (LOS) for surviving subjects was shorter at hospitals performing cardiac catheterization ($p < 0.01$). After adjusting for all variables, there were no independent associations between survival or LOS and hospital characteristics.

Conclusions: Some subsets of hospitals displayed higher survival and shorter LOS for OOHCA subjects but there was no independent association between hospital characteristics and outcome.

© 2010 Elsevier Ireland Ltd. All rights reserved.

Out-of-hospital cardiac arrest (OOHCA) has an incidence of 52.1 per 100,000 population, making it the third-leading cause of death in North America¹. Survival after OOHCA treated by emergency medical services (EMS) is about 7.9%, but varies greatly between

sites and regions (median 8.4%, ranging between 3.0% and 16.3%).¹ This large variation in survival is associated with differences in the characteristics of both patients, circumstances of arrest, and with EMS or hospital systems of care. The relative contributions of these factors to survival are unknown.

One possible influence on patient survival is the level of care received in the hospital after successful resuscitation by EMS. For example, specific interventions such as hypothermia^{2,3} and multidisciplinary changes in hospital care^{4,5} produced twofold increases in survival after OOHCA. Several papers have suggested the existence of hospital-related differences in survival after OOHCA.^{6–9} Observational studies suggest that early percutaneous coronary intervention can improve outcomes after cardiac arrest.^{10,11} However, the association between spe-

[☆] A Spanish translated version of the abstract of this article appears as Appendix in the online version at doi:10.1016/j.resuscitation.2009.12.006.

^{☆☆} A preliminary version of these data was presented at the Resuscitation Science Symposium, New Orleans, LA, November 5, 2008, and appear in abstract form in the proceedings.³⁷

* Corresponding author at: University of Pittsburgh, Department of Emergency Medicine, 3600 Forbes Ave #400A, Pittsburgh, PA 15261, United States.
Tel.: +1 412 647 9047; fax: +1 412 647 6999.

E-mail address: callawaycw@upmc.edu (C.W. Callaway).

cific characteristics of hospitals and survival after OOHCA is unknown.

Most hospitals treat few post-cardiac arrest patients.⁶ Therefore, it is difficult to compare individual hospital survival rates. However, in a large database, survival can be compared between categories of hospitals. The Resuscitations Outcome Consortium (ROC) developed and implemented a prospective, EMS-based registry to describe processes of care and outcome of patients assessed by EMS for OOHCA.¹² The present study used this database to test the hypothesis that survival to hospital discharge was associated with characteristics of receiving hospitals, including size (bed number), capability of performing cardiac catheterization, and total volume of OOHCA patients received.

1. Methods

Data were obtained from the ROC Cardiac Arrest Epidemiological Registry or “Epistry” which has been described in detail previously.¹² This registry is a prospective database of all persons evaluated for OOHCA by participating EMS agencies. Over 250 EMS agencies at sites in Canada and the US contribute data on an ongoing basis. Multiple mechanisms for surveillance are used at local sites to assure investigators of capture of all appropriate cases. Data are abstracted from EMS records and hospital records by data coordinators who use common data definitions. Data are transmitted to a central coordinating center by web-entry into an electronic database or batch upload of multiple cases grouped together. Within-form error checks are used to decrease data entry errors.

Subjects included were OOHCA patients of any age who received hospital care. “Cardiac arrest” was defined as receiving either (1) chest compressions by professional responder (paramedic, first-responder or other health professional arriving as part of the organized EMS response), or (2) rescue shock by professional responder or a bystander using an automated external defibrillator (AED). Cardiac arrest cases that were associated with trauma were excluded. “Received hospital care” was defined as either (1) arrived at the hospital with pulses, or (2) arrived at the hospital without documented pulses, but was discharged or died > 1 day later (i.e. was not pronounced dead on arrival at the hospital). When the subject was delivered to one hospital, and transferred to a second hospital on the same day, the second hospital was considered to be the treating hospital. These definitions were selected in order to exclude cases of EMS-treated OOHCA who never had restoration of pulses, and therefore who were not exposed to in-hospital post-resuscitation care. Subjects were entered into the database between December 1, 2005 and July 1, 2007, a time-period that excludes any interventional ROC trials.

Hospital characteristics were obtained from three sources. First, the EMS Structures Survey collected standardized data from each

ROC site completed by local investigators who described the local hospitals that receive patients from their EMS systems.¹³ Second, the American Hospital Directory is a publicly available database on US hospitals that is collated from a variety of public and payor databases (Medicare, licensed beds) (www.ahd.com). Third, the 2006–2007 Guide to Canadian Healthcare Facilities is a collation of data on Canadian hospitals provided by member hospitals (vol. 14, 2006–2007, www.cha.ca). Investigators examined the data from all three sources for each hospital that received subjects in the Epistry database in order to collate hospital characteristics. In the event of discrepancies, the report of local investigators who were familiar with the local hospital was used.

Hospitals were categorized based on several characteristics. Number of beds was used to categorize hospitals as large (>400 beds), medium (251–400 beds), and small (<250 beds). Capacity to perform acute cardiovascular interventions was defined as the presence of a cardiac catheterization laboratory, but the databases did not distinguish between capacity to perform emergent and elective catheterization or PCI. Trauma center designation was categorized as Level 1, Level 2, or non-trauma center. A teaching hospital was defined as one that listed active residency programs based at that hospital.

Patient and arrest characteristics were tabulated with descriptive statistics and compared between the number of hospital beds (size category) and cardiac catheterization capabilities of hospitals. Sample sizes within cardiac catheterization designation were similar across the different hospital size categories. Chi-square tests examined the independent association of hospital size category and catheterization capability with the primary outcome, survival to hospital discharge. Hospital length of stay was considered as a secondary outcome.

To account for known potential confounders, we used multiple logistic regression to examine the association of hospital type and survival to hospital discharge. We adjusted for variables previously associated with outcome: site; hospital characteristics including trauma level designation (1, 2, non-trauma), teaching institution; patient characteristics including gender, age (<1, 1–11, 12–19, 20–39, 40–60, 61–75, >75, unknown), witnessed collapse (witnessed, unwitnessed, or unknown), initial ECG rhythm (VT/VF, PEA, asystole, AED no-shock, or cannot determine), bystander attempts at cardiopulmonary resuscitation (CPR), and EMS process characteristics including response time (time from call to dispatch to first EMS vehicle arrival at scene). Variables were considered significantly associated with the outcome if $p < 0.05$.

In order to describe the relationship between hospital volume of OOHCA patients and outcome, we plotted survival vs. annualized number of cardiac arrests in the database for all hospitals. To avoid bias against hospitals with small volumes of cardiac arrest patients (<10 post-cardiac arrest patients a year), this analysis included all hospitals, regardless of size.

Table 1

Study hospital characteristics between US and Canadian ROC sites (* $p < 0.05$ between US and Canadian hospitals).

	All hospitals (n = 254)	Canadian hospitals (n = 110)	US hospitals (n = 144)
Number of beds			
≤100	42 (16.5%)	21 (19.1%)	21 (14.6%)
101–200	60 (23.6%)	19 (17.3%)	41 (28.5%)
201–300	49 (19.3%)	16 (14.5%)	33 (22.9%)
301–400	32 (12.6%)	16 (14.5%)	16 (11.1%)
>400	46 (18.1%)	24 (21.8%)	22 (15.3%)
Missing bed number	25 (9.8%)	14 (12.7%)	11 (7.6%)
Catheterization laboratory	140 (55.1%)	27 (24.5%)	113 (78.5%)*
Electrophysiology laboratory	38 (15.0%)	17 (16%)	21 (15%)
Trauma level 1	36 (14%)	17 (16%)	19 (13%)
Trauma level 2	19 (8%)	10 (9%)	9 (6%)
Residency program	111 (44%)	33 (30%)	78 (54%)*

Analyses of data were performed in S-Plus, v 6.2 (TIBCO Software, Palo Alto CA) or STATA v9.1 (StataCorp LP, College Station TX).

2. Results

Complete data were available from 8 sites in the US and 2 in Canada. EMS agencies delivered subjects to a total of 254 hospitals (144 in US and 110 in Canada). The distribution of hospital sizes was similar in the US and Canada (Table 1). A larger proportion of US hospitals performed cardiac catheterization and maintained residency programs than Canadian hospitals (Table 1). About 55% of hospitals reported being able to perform cardiac catheterization. Similar proportions of hospitals were designated Level 1 or 2 Trauma Centers in both US and Canada.

During this time interval, EMS treated 28,974 OOHCA cases and transported 10,496 cases to a hospital. Among these, 163 subjects (1.6%) were transported to hospitals outside of the surveillance area of the database, and no further information is available for those subjects. Of the remaining 10,333 subjects, 2395 (22.8%) had pulses at ED arrival while an additional 1767 (16.8%) did not have a pulse upon ED arrival or lacked EMS documentation of a pulse, but died or were discharged ≥ 1 days later. Survival data or other key data were missing for 75 cases (1.8%), leaving a total of 4087 subjects who received hospital care after OOHCA. Overall survival to hospital discharge for the 4087 patients who received hospital care was 32%.

A larger proportion of subjects delivered to the hospital with pulses or hospitalized >1 day were treated at large (37%) vs. medium (29%) and small (29%) hospitals (Table 2). Overall, 68% of subjects were treated at hospitals with cardiac catheterization capabilities. Hospitalized subjects were more often male and had an initial ECG rhythm of ventricular fibrillation or ventricular tachycardia (VF/VT) (Table 3). Median length of stay was longer in patients who survived to discharge (Table 4). For surviving patients, length of stay was longer at hospitals without cardiac catheterization capabilities ($p < 0.01$). This association was confounded by the fact that more Canadian hospitals, which have different incentive for length of stay, lacked cardiac catheterization capability. In a multivariable regression adjusting for country, initial VF/VT rhythm, age and witnessed collapse, the length of stay was not significantly associated with cardiac catheterization capability (0.18, 95% CI $-1.47, 1.82$), medium size (0.41, 95% CI $-1.12, 1.95$) or large size (0.32, 95% CI $-1.28, 1.92$).

Survival to hospital discharge had a significant univariable relationship with the availability of cardiac catheterization (OR = 1.43 95% CI 1.26–1.61, $p < 0.001$). Survival to hospital discharge was greater in hospitals with cardiac catheterization capability than those without (34% vs. 29%) (Fig. 1). This association was more prominent in small (<250 bed) hospitals (36% vs. 27%), than in medium (34% vs. 30%) or large hospitals (33% vs. 25%). Number of hospital beds did not have a significant relationship with survival ($p = 0.67$).

Subject characteristics associated with survival in the multivariable logistic model included an initial rhythm of VF/VT, younger age, and witnessed collapse ($p < 0.001$). After adjusting for all covariates, the odds of death in a hospital with cardiac catheterization capabilities vs. a hospital without cardiac catheterization was 0.79 (95% CI 0.62–1.001). The odds of death relative to a small hospital was 0.92 (95% CI 0.74–1.15) for medium-sized hospitals and 1.05 (95% CI 0.83–1.34) for large hospitals.

Survival rates did vary by volume of cardiac arrest cases treated (Table 5) with a high of 37.3% in hospitals with an annualized enrollment of ≥ 40 ROC patients a year and a low of 28.3% in hospitals with an annualized enrollment of 30–39 ROC patients a year. Survival




Fig. 1. Survival to discharge was greater in CATH than in no CATH hospitals regardless of bed number. Odds of death (triangles = No CATH; squares = CATH) were adjusted for witnessed collapse, VF/VT, and age.

was associated with case volume (Chi-square = 9.8, $p = 0.04$), but this relationship was attenuated when odds of death were adjusted for all covariates (Table 5).

3. Discussion

After resuscitation by EMS or in the ED after OOHCA, the unadjusted probability of survival to discharge is associated with care at a hospital capable of cardiac catheterization or treating ≥ 40 OOHCA patients annually. Mean length of stay is also shorter for patients hospitalized at hospitals capable of cardiac catheterization. The characteristics of patients delivered to different categories of hospitals do not differ, but after adjusting for response and patient characteristics, there was no independent association between survival and hospital characteristics. Therefore, it remains unclear whether EMS selection results in preferential transport of patients more or less likely to survive to specific hospitals. These data suggest some predilection for the most viable post-cardiac arrest patients to be treated at hospitals capable of cardiac catheterization.

The principal limitation of this study is that we are unable to describe whether therapeutic hypothermia, cardiac catheterization or other procedures were actually applied to individual patients. Cardiac catheterization availability may be surrogate for a more comprehensive cardiac and critical care program. Future studies should examine the influence of specific in-hospital procedures and therapies. Additional analyses with neurological and functional outcomes were not attempted because the current database lacks

Table 2

Distribution of subjects and hospitals defined by number of beds and cardiac catheterization (CATH) availability.

	Small (≤ 250 beds)	Medium (251–400 beds)	Large (>400 beds)	Missing ^a	Total
CATH					
Subjects (N)	963	790	897	137	2787
Hospitals (N)	50	30	31	6	117
No CATH					
Subjects (N)	529	396	277	98	1300
Hospitals (N)	51	17	12	6	86
Missing^a					
Subjects (N)	0	0	0	0	0
Hospitals (N)	0	0	0	0	0
Total					
Subjects (N)	1492	1186	1174	235	4087
Hospitals (N)	101	47	43	12	203 ^b

^a Some subjects were transferred to hospitals for which bed number ($n = 235$ subjects in 12 hospitals) were not known.^b A subset of 51 hospitals treated no subjects during the study interval.**Table 3**

Characteristics of subjects vs. hospital bed number and cardiac catheterization (CATH) capability. Designations are the same as in Table 2.

	Total N = 4087	Large CATH N = 897	Medium CATH N = 790	Small CATH N = 963	Large No CATH N = 277	Med No CATH N = 396	Small No CATH N = 529
Median age in years (IQR)	66 (25)	65 (27)	62 (23)	65 (24)	71 (22)	69 (24)	68 (22)
%Male	63%	62%	64%	60%	62%	64%	64%
Median Interval Dispatch to EMS Arrival in min (IQR)	5.0 (3.0)	5.0 (3.0)	5.0 (2.0)	5.0 (2.0)	6.0 (2.0)	5.0 (3.0)	6.0 (3.0)
Initial ECG rhythm							
VF/VT	41%	40%	37%	41%	44%	44%	40%
PEA	23%	21%	26%	27%	14%	16%	24%
Asystole	22%	21%	24%	22%	28%	23%	19%
Witnessed collapse							
EMS	14%	13%	14%	13%	16%	12%	13%
Bystander	51%	51%	48%	50%	55%	54%	52%
None	29%	26%	28%	28%	29%	33%	31%

Table 4

Hospital length of stay by hospital bed number and cardiac catheterization (CATH) capability.

	Total N = 4087	Large CATH N = 897	Medium CATH N = 790	Small CATH N = 963	Large No CATH N = 277	Medium No CATH N = 396	Small No CATH N = 529
Length of stay—all subjects median (IQR) days	3.0 (8.0)	3.0 (8.0)	4.0 (8.0)	3.0 (7.0)	2.0 (7.0)	3.0 (8.3)	2.5 (7.0)
Length of stay—discharged alive median (IQR) days	10.0 (12.0)	11.0 (13.8)	9.0 (9.0)	8.0 (9.0)	16.0 (20.0)	15.5 (27.8)	12.0 (13.0)
Length of stay—died in-hospital median (IQR) days	1.0 (3.0)	2.0 (3.0)	2.0 (4.0)	1.0 (3.0)	1.0 (3.0)	1.0 (3.0)	1.0 (3.0)

complete information on these measures at the time of discharge. One advantage of this prospective database is that future data collection can be expanded to record the actual cardiac procedures and treatment for individual subjects as well as more detailed outcomes. Those data will better answer what specific types of hospital care are associated with survival and improved outcome.

This study is also limited by the multiple out-of-hospital factors that affect patient outcome. When adjusted for all available clinical predictors, there was no significant independent effect of cardiac catheterization availability on survival. This result is similar to a recent analysis of survival after in-hospital cardiac arrest.¹⁴ In that analysis, care at a tertiary care facility resulted in higher unad-

Table 5Outcomes by hospital volume of OOHCA cases. Hospitals with no treated cases ($n = 51$) are excluded.

Yearly ROC cases	Number of hospitals	Survival to discharge (%)	Adjusted odds of death (95% CI)*
1–9	103	28.7%	Reference
10–19	55	30.8%	0.85 (0.65, 1.12)
20–29	23	32.6%	0.89 (0.65, 1.21)
30–39	11	28.3%	1.04 (0.75, 1.45)
≥ 40	11	37.3%	0.91 (0.67, 1.25)

* Adjusted for witnessed collapse, VF/VT, age and cardiac catheterization capability.

justed survival rate, but this effect was diminished when adjusted for differences between the hospital populations. In another series, differences in survival after OOHCA between two hospital systems were attributable in large part to out-of-hospital factors.¹⁵ Those prior studies and the present data underline the importance of optimizing multiple aspects of care, and not relying on a single aspect of the entire system of treatment, both in-hospital and out-of-hospital, in order to improve survival after cardiac arrest.

Availability of cardiac catheterization was expected to be beneficial because of the high incidence of coronary artery occlusion in out-of-hospital cardiac arrest.^{16,17} Primary percutaneous coronary intervention (PCI) by experienced cardiologists is associated with superior survival and fewer complications for patients with ST-elevation myocardial infarction (STEMI),¹⁸ and also may benefit non-STEMI patients.^{19–21} This study did not directly compare early intervention to conservative management, but do indicate that post-cardiac arrest patients are currently directed towards hospitals with availability of coronary intervention.

Invasive coronary interventions may be under-utilized for post-cardiac arrest patients.²² Some authors have proposed triage and transfer of high-risk cardiac patients without cardiac arrest to cardiac centers, which are capable of primary PCI,²³ and some regions also transfer patients after cardiac arrest.²⁴ The benefit of transfer and longer transport may only affect outcomes for selected “high-risk” STEMI patients,²⁵ and emergent PCI may only benefit a subset of post-cardiac arrest patients.²⁶ Regardless, more data are required to establish whether treatment of post-cardiac arrest patients in a fashion similar to high-risk STEMI patients is beneficial.

This study adds to the discussion about whether patients with restoration of pulses after OOHCA should be transported to the nearest hospital or to specialized centers.²⁷ In many critical illnesses, specialized care improves outcomes.²⁸ For example, implementation of trauma systems is associated with improved survival after life-threatening injury,^{29,30} particularly at individual trauma centers with higher volumes of cases.³¹ Likewise, care of stroke patients at specialized stroke units is associated with better survival and better functional recovery, independent of reperfusion.^{32,33} Better survival in these studies was related to improved systems of care rather than single interventions. This paper helps identify the important covariates that should be measured in order to test whether similar systems of care also may benefit OOHCA.

The present data suggest that patients with greater chance of survival are being treated at larger cardiac hospitals, but cannot separate the contribution of hospital and patient characteristics. Other retrospective analyses found relationships between survival after OOHCA and different aspects of hospital care, including case volume and coronary interventions.^{6,7,34–36} The present analysis provided a more complete description than a previous study⁶ that excluded hospitals with less than 12 patients a year, and found no independent effect of case volume. One limitation of the present analysis is the fact that some hospitals would treat post-cardiac arrest patients delivered to the hospital from sources other than ROC EMS agencies. For most facilities, the ROC database does include the majority of EMS services serving the surrounding geographical region. Thus, the volume of cardiac arrest patients used in this analysis is probably a good surrogate for total volume of post-cardiac arrest patients.

In summary, patients with restoration of pulses after out-of-hospital cardiac arrest who were treated at hospitals capable of invasive cardiac procedures and hospitals that treat a large volume of cardiac arrest patients had increased rates of survival. However, there was no independent effects of hospital characteristics after adjusting for multiple other factors associated with survival after cardiac arrest. These associations underline the need to study implementation of regional systems of care for post-

cardiac arrest patients in a prospective fashion with attention to multiple important covariates. Future research should determine if these observations are robust and identify whether specific hospital procedures are associated with survival.

Conflict of interest

The authors do not have any direct conflicts of interest related to the topics in this paper.

Acknowledgements

Funding sources: The ROC is supported by a series of cooperative agreements to 10 regional clinical centers and one Data Coordinating Center (5U01 HL077863, HL077881, HL077871 HL077872, HL077866, HL077908, HL077867, HL077885, HL077887, HL077873, HL077865) from the National Heart, Lung and Blood Institute in partnership with the National Institute of Neurological Disorders and Stroke, U.S. Army Medical Research & Materiel Command, The Canadian Institutes of Health Research (CIHR)—Institute of Circulatory and Respiratory Health, Defence Research, the American Heart Association and Development Canada and the Heart and Stroke Foundation of Canada.

References

- Nichol G, Thomas E, Callaway CW, et al. Resuscitation Outcomes Consortium Investigators. Regional variation in out-of-hospital cardiac arrest incidence and outcome. *JAMA* 2008;300:1423–31.
- Hypothermia after Cardiac Arrest Study Group. Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. *N Engl J Med* 2002;346:549–56.
- Bernard SA, Gray TW, Buist MD, et al. Treatment of comatose survivors of out-of-hospital cardiac arrest with induced hypothermia. *N Engl J Med* 2002;346:557–63.
- Sunde K, Pytte M, Jacobsen D, et al. Implementation of a standardised treatment protocol for post resuscitation care after out-of-hospital cardiac arrest. *Resuscitation* 2007;73:29–39.
- Rittenberger JC, Guyette FX, Tisherman SA, DeVita MA, Alvarez RJ, Callaway CW. Outcomes of a hospital-wide plan to improve care of comatose survivors of cardiac arrest. *Resuscitation* 2008;79:198–204.
- Carr BG, Kahn JM, Merchant RM, Kramer AA, Neumar RW. Inter-hospital variability in post-cardiac arrest mortality. *Resuscitation* 2009;80:30–4.
- Langhelle A, Tyvoll SS, Lexow K, Hapnes SA, Sunde K, Steen PA. In-hospital factors associated with improved outcome after out-of-hospital cardiac arrest. A comparison between four regions in Norway. *Resuscitation* 2003;56:247–63.
- Skrifvars MB, Pettilä V, Rosenberg PH, Castrén M. A multiple logistic regression analysis of in-hospital factors related to survival at six months in patients resuscitated from out-of-hospital ventricular fibrillation. *Resuscitation* 2003;59:319–28.
- Liu JM, Yang Q, Pirralo RG, Klein JP, Aufderheide TP. Hospital variability of out-of-hospital cardiac arrest survival. *Prehosp Emerg Care* 2008;12:339–46.
- Peels HO, Jessurun GAJ, van der Horst ICC, Arnold AER, Piers LH, Zijlstra F. Outcome in transferred and nontransferred patients after percutaneous coronary intervention for ischaemic out-of-hospital cardiac arrest. *Cathet Cardiovasc Interv* 2008;71:147–51.
- Garot P, Lefevre T, Eltchaninoff H, et al. Six-month outcome of emergency percutaneous coronary intervention in resuscitated patients after cardiac arrest complicating ST-elevation myocardial infarction. *Circulation* 2007;115:1354–62.
- Morrison LJ, Nichol G, Rea TD, et al., ROC Investigators. Rationale, development and implementation of the Resuscitation Outcomes Consortium Epistry-Cardiac Arrest. *Resuscitation* 2008;78:161–9.
- Davis DP, Garberson IA, Andrusiek DL, et al. A descriptive analysis of Emergency Medical Service Systems participating in the Resuscitation Outcomes Consortium (ROC) network. *Prehosp Emerg Care* 2007;11:369–82.
- Skrifvars MB, Castrén M, Aune S, Thoren AB, Nurmi J, Herlitz J. Variability in survival after in-hospital cardiac arrest depending on the hospital level of care. *Resuscitation* 2007;73:73–81.
- Hollenberg J, Lindqvist J, Ringh M, et al. An evaluation of post-resuscitation care as a possible explanation of a difference in survival after out-of-hospital cardiac arrest. *Resuscitation* 2000;74:242–52.
- Spaulding CM, Joly LM, Rosenberg A, et al. Immediate coronary angiography in survivors of out-of-hospital cardiac arrest. *N Engl J Med* 1997;336:1629–33.

17. Reynolds JC, Callaway CW, El Khoudary SR, Moore CG, Alvarez RJ, Rittenberger JC. Coronary angiography predicts improved outcome following cardiac arrest: propensity-adjusted analysis. *J Intens Care Med* 2009;24:179–86.
18. Antman EM, Hand M, Armstrong PW, et al. 2007, Focused Update of the ACC/AHA 2004 Guidelines for the management of patients with ST-elevation myocardial infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines: developed in collaboration With the Canadian Cardiovascular Society endorsed by the American Academy of Family Physicians: 2007 Writing Group to Review New Evidence and Update the ACC/AHA 2004 Guidelines for the Management of Patients With ST-Elevation Myocardial Infarction, Writing on Behalf of the 2004 Writing Committee. *Circulation* 2008;117:296–329.
19. Choudhry NK, Singh JM, Barolet A, Tomlinson GA, Detsky AS. How should patients with unstable angina and non-ST-segment elevation myocardial infarction be managed? A meta-analysis of randomized trials. *Am J Med* 2005;118:465–74.
20. Hoenig MR, Doust JA, Aroney CN, Scott IA. Early invasive versus conservative strategies for unstable angina & non-ST-elevation myocardial infarction in the stent era. *Cochrane Database Syst Rev* 2006;3. CD004815.
21. O'Donoghue M, Boden WE, Braunwald E, et al. Early invasive vs conservative treatment strategies in women and men with unstable angina and non-ST-segment elevation myocardial infarction: a meta-analysis. *JAMA* 2008;300:71–80.
22. Merchant RM, Abella BS, Khan M, et al. Cardiac catheterization is underutilized after in-hospital cardiac arrest. *Resuscitation* 2008;79:398–403.
23. Henry TD, Atkins JM, Cunningham MS, et al. ST-segment elevation myocardial infarction: recommendations on triage of patients to heart attack centers: is it time for a national policy for the treatment of ST-segment elevation myocardial infarction? *J Am Coll Cardiol* 2006;47:1339–45.
24. Henry TD, Sharkey SW, Burke MN, et al. A regional system to provide timely access to percutaneous coronary intervention for ST-elevation myocardial infarction. *Circulation* 2007;116:721–8.
25. Rathore SS, Epstein AJ, Nallamothu BK, Krumholz HM. Regionalization of ST-segment elevation acute coronary syndromes care: putting a national policy in proper perspective. *J Am Coll Cardiol* 2006;47:1346–9.
26. Anyfantakis ZA, Baron G, Aubry P, et al. Acute coronary angiographic findings in survivors of out-of-hospital cardiac arrest. *Am Heart J* 2009;157:312–8.
27. Lurie KG, Idris A, Holcomb JB. Level 1 cardiac arrest centers: learning from the trauma surgeons. *Acad Emerg Med* 2005;12:79–80.
28. Pronovost PJ, Angus DC, Dorman T, Robinson KA, Dremsizov TT, Young TL. Physician staffing patterns and clinical outcomes in critically ill patients: a systematic review. *JAMA* 2002;288:2151–62.
29. Nathens AB, Jurkovich GJ, Maier RV, et al. Relationship between trauma center volume and outcomes. *JAMA* 2001;285:1164–71.
30. Celso B, Tepas J, Langland-Orban B, et al. A systematic review and meta-analysis comparing outcome of severely injured patients treated in trauma centers following the establishment of trauma systems. *J Trauma* 2006;60:371–8.
31. MacKenzie EJ, Rivara FP, Jurkovich GJ, et al. A national evaluation of the effect of trauma-center care on mortality. *N Engl J Med* 2006;354:366–78.
32. Foley N, Salter K, Teasell R. Specialized stroke services: a meta-analysis comparing three models of care. *Cerebrovasc Dis* 2007;23:194–202.
33. Stroke Unit Trialists' Collaboration. Organised inpatient (stroke unit) care for stroke. *Cochrane Database Syst Rev* 2007;4:CD000197.
34. Herlitz J, Engdahl J, Svensson L, Angquist KA, Silfverstolpe J, Holmberg S. Major differences in 1-month survival between hospitals in Sweden among initial survivors of out-of-hospital cardiac arrest. *Resuscitation* 2006;70:404–9.
35. Herlitz J, Bang A, Gunnarsson J, et al. Factors associated with survival to hospital discharge among patients hospitalised alive after out of hospital cardiac arrest: change in outcome over 20 years in the community of Göteborg, Sweden. *Heart* 2003;89:25–30.
36. Werling M, Thorén AB, Axelsson C, Herlitz J. Treatment and outcome in post-resuscitation care after out-of-hospital cardiac arrest when a modern therapeutic approach was introduced. *Resuscitation* 2007;73:40–5.
37. Callaway CW, Schmicker R, Kampmeyer M, et al. Influence of receiving hospital characteristics on survival after cardiac arrest. *Circulation* 2008;118:S1446.