

# Acid–base physiology: comments on 10 contentious assertions

Thomas J Morgan, Balasubramanian Venkatesh and Rinaldo Bellomo

In the world of acid–base, models to explain observed phenomena tend to be adopted by particular “schools”, and then fiercely defended, on an almost tribal basis.<sup>1</sup> Meanwhile, irrefutable truth (as distinct from debating positions) can remain elusive. In this context, we offer brief comments on 10 assertions which have had or still have currency. We regard these responses as consistent with available evidence, but acknowledge that all may require modification or even abandonment, subject to further discoveries.

## 1. The Stewart approach is sufficient to explain acid–base physiology

In the Stewart model,<sup>2,3</sup> three independent variables ( $P_{CO_2}$ , the total concentration of non-volatile weak acid (Atot) and the strong ion difference [SID]) directly control plasma concentrations of several dependent variables, including pH and  $HCO_3^-$ . Six simultaneous equations (Table 1), separately or as a fourth-order polynomial, shed light on many areas of acid–base physiology (see below), but with several caveats. First, some prefer to characterise strong ions as mere “spectator” ions, their net charge a convenient inverse quantification of the “buffer base” concentration, which is the total concentration of bicarbonate and non-bicarbonate buffers.<sup>4-6</sup> This more traditional approach designates the buffer base as the site of all relevant molecular acid–base transactions. Stewart advocates have also been criticised for treating water as a limitless proton “reservoir” (Figure 1), ignoring the alternative viewpoint that it acts more as a proton conduit via excited-state “water bridges”.<sup>8</sup> Hence, for some, the Stewart approach, while mathematically accurate, is mechanistically incorrect.<sup>5,9</sup> The debate continues.

**Table 1. The six equations of Stewart\***

- $[H^+] \times [OH^-] = K'_w$
- $[H^+] \times [A^-] = K_a \times HA$
- $[HA] + [A^-] = Atot$
- $[H^+] \times [HCO_3^-] = K_c \times P_{CO_2}$
- $[H^+] \times [CO_3^{2-}] = K_d \times [HCO_3^-]$
- $SID + [H^+] - [HCO_3^-] - [CO_3^{2-}] - [A^-] - [OH^-] = 0$

SID = strong ion difference (net charge concentration of fully dissociated ions). Atot = total concentration of non-volatile weak acids. \* All K values are known dissociation constants.


There is one genuine discrepancy in Stewart’s original concept. Modelling plasma in isolation disregards ion and water traffic and associated Gibbs–Donnan phenomena in the interstitium–plasma–erythrocyte (IPE) complex.<sup>10</sup> Because of this, independent status cannot be claimed for plasma SID in vivo, since  $CO_2$ -driven traffic causes it to alter with  $P_{CO_2}$ .<sup>3,11</sup>  $P_{CO_2}$  independence is retained by IPE SID, a more nebulous entity.

## 2. Acidosis stimulates renal bicarbonate reabsorption, new bicarbonate generation and proton excretion

These traditional concepts can be usefully reframed in Stewart terms as follows:<sup>3,12,13</sup>

- Tubular proton and bicarbonate transporters are simultaneous strong ion transporters, directly or indirectly, with immediate effects on urinary SID.
- At low urinary pH,  $NH_4^+$  (pKa, 9.2) is a strong cation. However, in extracellular fluid,  $NH_4^+$  is a weak cation, present in minute concentrations. It is therefore excluded from the urinary SID calculation.

**Figure 1. The pH of pure water v temperature\***



\* Graph based on data from Harned and Owen,<sup>7</sup> showing that the pH of a water solution can be changed without adding any “acid”. The implications are that water itself is a source of interactive hydrogen ions, and that it is impossible to identify the source of hydrogen ions with certainty in aqueous solutions such as plasma.

## POINTS OF VIEW

- Renal ionophore responses to acidosis decrease urinary SID.
- Upregulation of tubular  $\text{NH}_3$  synthesis in acidosis supports urinary SID reduction, substituting  $\text{NH}_4^+$  for sodium as a cationic partner of chloride.
- Decreasing urinary SID increases extracellular SID, restoring plasma pH and bicarbonate.

### 3. Hypoalbuminaemia does not affect acid–base physiology

Albumin acts as a weak acid.<sup>2,3</sup> Experiments using magnetic resonance and confirmed by modelling<sup>14</sup> quantify the titratable protein charge in plasma at physiological pH at 12–13 mEq/L,<sup>15</sup> mostly from histidine moieties on albumin.<sup>16</sup> Hypoalbuminaemia diminishes these acidifying effects and blood is alkalinised. The net negative charge of albumin necessitates an “albumin-corrected” anion gap to counteract albumin fluctuations.<sup>16</sup>

### 4. Acidosis in shock is from proton release generated by ATP hydrolysis<sup>17</sup>

No studies of clinical shock show proton release from ATP hydrolysis. In fact experimentally, net ATP hydrolysis implies zero perfusion (death or full ischaemia).<sup>18</sup> ATP depletion has never been shown in clinical septic, haemorrhagic, anaphylactic or cardiogenic shock or by magnetic resonance spectroscopy in extreme experimental septic shock.<sup>19</sup> A more elegant explanation is that tissues release L-lactate under physiological stress via adrenergically mediated accelerated glycolysis,<sup>20</sup> reducing SID and hence acidifying plasma.

### 5. Hyperlactataemia implies tissue hypoxia

The concept that tissue hypoxia underlies most hyperlactaemic states lacks any empirical basis.<sup>21</sup> With no gold standard to diagnose tissue hypoxia, its presence in septic shock, for example, can be challenged.<sup>22</sup> During maximal exercise, plasma lactate increases despite preserved intracellular oxygenation,<sup>23</sup> whereas in extreme hypoxaemia (eg, a person on Mount Everest) lactate is, at most, minimally elevated.<sup>22</sup> Lactate shuttles within cells and among tissues as a major biofuel.<sup>23</sup> Although it is an independent mortality predictor in shock, evidence suggests that hyperlactataemia is due to accelerated aerobic glycolysis driven by adrenergic stress,<sup>24</sup> facilitating bioenergetic efficiency in muscle, heart and brain cells. These characteristics best fit the notion of an adaptive survival response.

### 6. Lactate-balanced solutions contain “bicarbonate as lactate”

There is a belief that administered lactate is oxidised to bicarbonate, forming the rationale for “lactate buffered”

solutions. In reality, manufacturers substitute organic anions for bicarbonate to avert bicarbonate degradation by  $\text{CO}_2$  loss on autoclaving and storage. Administered L-lactate is metabolised to pyruvate and then predominantly to glucose (gluconeogenesis) and  $\text{CO}_2$  (via the Krebs cycle). Its rapid metabolic “disappearance” means that the effective SID of a lactate-balanced solution is the difference between non-metabolised cations and anions. The effective SID of Hartmann’s solution is therefore 29 mEq/L, slightly exceeding that required for exact balance (24 mEq/L).<sup>25</sup>

### 7. The bicarbonate in sodium bicarbonate corrects metabolic acidosis

In Stewart terms, the critical property of sodium bicarbonate is the presence of sodium, a strong cation, without a corresponding strong anion such as chloride.<sup>26</sup> Its SID is therefore its sodium concentration (1000 mEq/L in a 1 M solution), which increases extracellular SID on administration and antagonises metabolic acidosis. The bicarbonate component is rapidly exhaled as  $\text{CO}_2$ . The SID of NaOH 1M is also 1000 mEq/L but with a pH of 14, it is unsuitable for intravenous use. Otherwise it would be equally effective in reversing metabolic acidosis, despite the absence of bicarbonate. Conversely, ammonium bicarbonate<sup>27</sup> contains no strong ions (its SID is zero). In sufficient quantity and despite the presence of bicarbonate, it will cause metabolic acidosis.<sup>25</sup>

### 8. Paradoxical intracellular acidosis from bicarbonate administration is a big problem

Tissue experiments confirm that the  $\text{CO}_2$  content of added sodium bicarbonate generates marked respiratory acidosis.<sup>28</sup> The in-vivo situation differs for the following reasons.<sup>26</sup>

- There is massive haemodilution by cardiac output.
- Any intravenous  $\text{CO}_2$  load immediately increases  $\text{CO}_2$  excretion via mixed venous pulmonary transit.
- Modelling indicates that during acidosis correction, only a fraction of administered  $\text{CO}_2$  in sodium bicarbonate requires subsequent excretion.<sup>26</sup>

In practice, large doses of sodium bicarbonate must be infused rapidly, for example, 1.5 mmol/kg in 5 minutes, to generate detectable  $\text{Paco}_2$  and  $\text{Vco}_2$  upswings.<sup>29</sup> Slower administration should prevent  $\text{CO}_2$  surges, with possible exceptions being extreme pulmonary hypoperfusion states.

### 9. Fluid-induced acidosis is simple bicarbonate dilution

Bicarbonate is  $\text{CO}_2$  in hydrated anionic form. Rapid  $\text{CO}_2$ – $\text{HCO}_3^-$  transitions are facilitated by carbonic anhydrase, and body fluids are open to  $\text{CO}_2$  ingress and egress. Hence, in living open systems during volume loading, there is no fixed bicarbonate mass to dilute. Concentrations continue to be

determined by  $P_{CO_2}$ ,  $Atot$  and  $SID$ .<sup>3</sup> Saline haemodilution reduces both  $SID$  and  $Atot$ , with the net result being a fall in bicarbonate.<sup>25</sup>

### 10. Loop diuretics cause metabolic alkalosis because of “volume contraction”

Loop diuretics decrease urinary  $SID$  by competing for the chloride site on the  $NaK_2Cl$  cotransporter in the thick ascending limb.<sup>30</sup> The output of reduced- $SID$  urine increases the extracellular  $SID$ , causing metabolic alkalosis. Although mineralocorticoid responses to diuretic-induced volume contraction also reduce urinary  $SID$ , they are readily overridden if the diuretic itself increases urinary  $SID$ , as does, for example, acetazolamide.<sup>31</sup>

### Competing interests

None declared.

### Author details

Thomas J Morgan, Senior Lecturer<sup>1</sup>

Balasubramanian Venkatesh, Professor of Intensive Care<sup>2</sup>

Rinaldo Bellomo, Professor of Intensive Care<sup>3</sup>

1 Mater Research Institute, University of Queensland, Brisbane, QLD, Australia.

2 Princess Alexandra and Wesley Hospitals, University of Queensland, Brisbane, QLD, Australia.

3 Australian and New Zealand Intensive Care Research Centre, Melbourne, VIC, Australia.

Correspondence: t.morgan@uq.edu.au

### References

- Severinghaus JW. Siggaard-Andersen and the “Great Trans-Atlantic Acid-Base Debate”. *Scand J Clin Lab Invest Suppl* 1993; 214: 99-104.
- Stewart PA. Modern quantitative acid-base chemistry. *Can J Physiol Pharmacol* 1983; 61: 1444-61.
- Morgan TJ. The Stewart approach— one clinician’s perspective. *Clin Biochem Rev* 2009; 30: 41-54.
- Singer RB, Hastings AB. An improved clinical method for the estimation of disturbances of the acid-base balance of human blood. *Medicine (Baltimore)* 1948; 27: 223-42.
- Kurtz I, Kraut J, Ornekian V, Nguyen MK. Acid-base analysis: a critique of the Stewart and bicarbonate-centered approaches. *Am J Physiol Renal Physiol* 2008; 294: F1009-31.
- Siggaard-Andersen O, Fogh-Andersen N. Base excess or buffer base (strong ion difference) as measure of a non-respiratory acid-base disturbance. *Acta Anaesthesiol Scand Suppl* 1995; 107: 123-8.
- Harned HS, Owen BB. The physical chemistry of electrolytic solutions. New York: Reinhold Publishing Corp, 1958: 634-49, 752-4.
- Gepshtein R, Leiderman P, Genosar L, Huppert D. Testing the three step excited state proton transfer model by the effect of an excess proton. *J Physical Chem A* 2005; 109: 9674-84.
- Doberer D, Funk GC, Kirchner K, Schneeweiss B. A critique of Stewart’s approach: the chemical mechanism of dilutional acidosis. *Intensive Care Med* 2009; 35: 2173-80.
- Wolf MB, Deland EC. A mathematical model of blood-interstitial acid-base balance: application to dilution acidosis and acid-base status. *J Appl Physiol* 2011; 110: 988-1002.
- Langer T, Scotti E, Carlesso E, et al. Electrolyte shifts across the artificial lung in patients on extracorporeal membrane oxygenation: interdependence between partial pressure of carbon dioxide and strong ion difference. *J Crit Care* 2015; 30: 2-6.
- Gattinoni L, Carlesso E, Cadringer P, Caironi P. Strong ion difference in urine: new perspectives in acid-base assessment. *Crit Care* 2006; 10: 137.
- Yunos NM, Bellomo R, Story D, Kellum J. Bench-to-bedside review: chloride in critical illness. *Crit Care* 2010; 14: 226.
- Watson PD. Modeling the effects of proteins on pH in plasma. *J Appl Physiol* (1985) 1999; 86: 1421-7.
- Staempfli HR, Constable PD. Experimental determination of net protein charge and  $A(tot)$  and  $K(a)$  of nonvolatile buffers in human plasma. *J Appl Physiol* (1985) 2003; 95: 620-30.
- Figge J, Mydosh T, Fencl V. Serum proteins and acid-base equilibria: a follow-up. *J Lab Clin Med* 1992; 120: 713-9.
- Hotchkiss RS, Karl IE. Reevaluation of the role of cellular hypoxia and bioenergetic failure in sepsis. *JAMA* 1992; 267: 1503-10.
- May C, Wan L, Williams J, et al. A technique for the measurement of renal ATP in a large animal model of septic shock. *Int J Artif Organs* 2005; 28: 16-21.
- May CN, Ishikawa K, Wan L, et al. Renal bioenergetics during early gram-negative mammalian sepsis and angiotensin II infusion. *Intensive Care Med* 2012; 38: 886-93.
- Levy B. Lactate and shock state: the metabolic view. *Curr Opin Crit Care* 2006; 12: 315-21.
- James JH, Luchette FA, McCarter FD, Fischer JE. Lactate is an unreliable indicator of tissue hypoxia in injury or sepsis. *Lancet* 1999; 354: 505-8.
- Garcia-Alvarez M, Marik P, Bellomo R. Sepsis-associated hyperlactatemia. *Crit Care* 2014; 18: 503.
- Garcia-Alvarez M, Marik P, Bellomo R. Stress hyperlactataemia: present understanding and controversy. *Lancet Diabetes Endocrinol* 2014; 2: 339-47.
- Levy B, Gibot S, Franck P, et al. Relation between muscle  $Na+K+ATPase$  activity and raised lactate concentrations in septic shock: a prospective study. *Lancet* 2005; 365: 871-5.
- Morgan TJ. The ideal crystalloid - what is ‘balanced’? *Curr Opin Crit Care* 2013; 19: 299-307.
- Morgan TJ. Buffers. In: Kellum JA, Elbers, PWG, editors. Stewart’s textbook of acid base. Amsterdam: AcidBase.org, 2009: 281-91.
- McKenzie DJ, Randall DJ, Lin H, Aota S. Effects of changes in plasma pH,  $CO_2$  and ammonia on ventilation in trout. *Fish Physiol Biochem* 1993; 10: 507-15.
- Levrout J, Giunti C, Ciebiera JP, et al. Initial effect of sodium bicarbonate on intracellular pH depends on the extracellular nonbicarbonate buffering capacity. *Crit Care Med* 2001; 29: 1033-9.
- Levrout J, Garcia P, Giunti C, et al. The increase in  $CO_2$  production induced by  $NaHCO_3$  depends on blood albumin and hemoglobin concentrations. *Intensive Care Med* 2000; 26: 558-64.
- Ellison DH. Diuretic drugs and the treatment of edema: from clinic to bench and back again. *Am J Kidney Dis* 1994; 23: 623-43.
- Moviat M, Pickkers P, van der Voort PH, van der Hoeven JG. Acetazolamide-mediated decrease in strong ion difference accounts for the correction of metabolic alkalosis in critically ill patients. *Crit Care* 2006; 10: R14. □